

Counting Extra Dimensions: Magnetic Cherenkov Radiation from High Energy Neutrinos

G. Domokos^{a1}, Andrea Erdas^{a,b2} and S. Kovesi-Domokos^{a1}

^a Department of Physics and Astronomy, The Johns Hopkins University,
Baltimore, MD

^b Dipartimento di Fisica dell'Università & INFN, Cagliari

Abstract In theories which require a space of dimension $d > 4$, there is a natural mechanism of suppressing neutrino masses: while Standard Model fields are confined to a 3-brane, right handed neutrinos live in the bulk. Due to Kaluza-Klein excitations, the effective magnetic moments of neutrinos are enhanced. The effective magnetic moment is a monotonically growing function of the energy of the neutrino: consequently, high energy neutrinos can emit observable amounts of magnetic Cherenkov radiation. By observing the energy dependence of the magnetic Cherenkov radiation, one may be able to determine the number of compactified dimensions.

1 Introduction

In theories with extra dimensions there is an interesting alternative to the conventional seesaw mechanism of giving neutrinos a small mass [1]. In that scheme, the left-handed neutrino (ν_L) resides on the brane together with all other Standard Model fields, whereas the right handed neutrinos (ν_R) live in the bulk. The neutrino mass of any flavor is then suppressed with respect to other fermion masses by a factor of the order of $\sim M_*/M_P$, M_P being the four dimensional Planck mass and M_* is the string scale. Its currently favored value is somewhere between ~ 10 TeV and ~ 100 TeV. Phenomenological implications of such a scheme have been studied by a number of authors. For our purposes, the relevant papers are the ones authored by Faraggi and Pospelov [2], McLaughlin and Ng [3] and by Agashe and Wu [4].

We argue that if indeed ν_R of any flavor lives in the bulk, then magnetic Cherenkov radiation by that neutrino flavor is enhanced due to the contribution of Kaluza-Klein (KK) excitations to the transition probability. In the next Section we outline the calculation leading to such a conclusion. We find that, in fact, there is hardly anything to calculate, since one is able to use results of ref. [3] combined with those of Sawyer [5]. In Sec. 3 practical aspects of the results are discussed;

¹e-mail: skd@jhu.edu

²e-mail: aerdas@pha.jhu.edu

in particular, we estimate the yield of Cherenkov photons in water, relevant for underwater detectors. (Estimates of yields in detectors in ice are somewhat lower, but not dramatically so.) The last Section is devoted to a discussion of the results.

It is to be noted that there exists an extensive literature on magnetic Cherenkov radiation of neutrinos: several results have been repeatedly rediscovered, rederived, *etc.* often with results which contradict each other. To our knowledge, the papers of Sawyer [5] and of Ionnisian and Raffelt [6] are internally consistent. In addition, ref. [6] contains an extensive list of references. Throughout this paper natural units are used: $\hbar = c = 1$.

2 Cherenkov radiation in the presence of KK excitations

If a particle is neutral, but of nonzero spin, it is scattered on an electromagnetic field due to its magnetic moment. By using the optical theorem and crossing symmetry, the magnitude of the magnetic moment can be extracted from the imaginary part of the forward (magnetic) Compton amplitude in the zero frequency limit. If the particle in question has excited states, this leads to an energy dependent *effective magnetic moment*: the imaginary part of the Compton amplitude depends on the number of excited states available at a given energy. This is, in fact, the argument put forward in ref. [3].

The forward Compton amplitude depends on the expression of the transition moments between the ground state of the particle and its excited states. Obviously, this reasoning is applicable to neutrinos with ν_R living in the bulk, due to the existence of KK excitations. Assuming toroidal compactification of the extra dimensions of equal radii ($R^{-1} = M_c$), it is obvious that the effective magnetic moment squared is proportional to the number of available KK states at a given energy, as well as to the mass squared of the neutrino:

$$\mu_{\text{eff}}^2 \propto m_D^2 \left(\frac{E}{M_c} \right)^n \quad (1)$$

McLaughlin and Ng [3] find for a reasonable choice of parameters, qualitatively consistent with solar and atmospheric neutrino data:

$$\mu_N \approx 1.6 \times 10^{-19} \mu_B \frac{m_D}{\text{1eV}}, \quad (2)$$

where μ_N is the transition moment between the zero mode of ν_L and the N^{th} KK level, while m_D stands for the (Dirac) mass of the neutrino. The quantity μ_B is the Bohr magneton, giving a convenient comparison scale. The result quoted above is valid in the weak mixing approximation, which is well justifiable here, see [2, 3]. Remarkably, the transition moment is independent of the KK level: this fact plays a significant role in what follows.

The resulting effective magnetic moment is given by:

$$\mu_{\text{eff}}^2 \approx 1.6 \times 10^{-15} \mu_B^2 \frac{\pi^{(1-n)/2}}{2^n \Gamma\left(\frac{n+1}{2}\right)} \left(\frac{E}{M_*} \right)^n, \quad (3)$$

where n stands for the number of compactified dimensions. The factor multiplying $(E/M_*)^n$ is proportional to the degeneracy factor of the KK states. In the last equation, the mass of the neutrino has been eliminated in favor of its Yukawa coupling to the Higgs field, y , according to the formula:

$$m_D^2 = y^2 V^2 \left(\frac{M_c}{2\pi M_*} \right)^n, \quad (4)$$

see ref. [1]. Here, V stands for the vacuum expectation value of the Higgs field; its standard value is being used. Due to our ignorance, we also set $y \approx 1$. As far as n , the number of compactified dimensions is concerned, certain models (superstring, SUGRA) suggest $n = 6$ or $n = 7$, respectively. However, it should be realized that at present, it is unclear whether we have a phenomenologically viable model describing physics beyond the Standard Model. For that reason, we prefer to keep an open mind about the number of compactified dimensions.

With present techniques, Cherenkov radiation can be detected typically in the visible and in the (near) UV. Hence, one can directly use the emission rates obtained by Sawyer [5], replacing $\mu^2 \rightarrow \mu_{\text{eff}}^2$. (This approximation is justified by the fact that the energy of the emitted Cherenkov photon is low compared to energy scales of the KK excitations.)

Using this, one obtains the formula for the number of photons emitted in a unit frequency interval and per unit length by a neutrino of $v \approx 1$:

$$\frac{dN}{d\omega dx} = \omega^2 (\mu_{\text{eff}})^2 (\epsilon(\omega) - 1). \quad (5)$$

It is amusing to observe that in the tree approximation used by Sawyer [5], the result of a lowest order perturbative calculation of the Cherenkov emission rate gives the same result as the old calculation of Ginzburg based on classical electrodynamics [7]. This is a fairly general result: the tree approximation to any given Feynman diagram can be expressed in terms of an iterative solution of the classical field equations to the given order. Taking into account KK excitations requires the use of a quantum calculation of the type devised by Sawyer, *l.c.* However, in the *effective moment approximation* as used above, one can still resort to the classical calculation.

3 Observability

Without the enhancement provided by the presence of KK excitations, it is hopeless to observe the magnetic Cherenkov radiation. Using the upper limits on the magnetic moments of neutrinos as given by the Particle Data Group [8], one finds that the number of photons emitted by a neutrino of any flavor is $\leq 10^{-16}/\text{km}$ or so. Clearly, the observation of such a rare event, let alone its identification as a Cherenkov emission is out of question in any detector at present or in the foreseeable future.

Do KK excitations help?

In order to answer that question, we assume a quantum efficiency of the PMTs in a detector to be around 10% and that about 100 photoelectrons have to be detected in a detector in order to identify a Cherenkov ring. Furthermore, we assume a detector size of the order of 1km. A simple model of phototube response consists of no response below a minimal wavelength, λ_{min} , in the UV, a flat response between λ_{min} and a fixed λ_{max} , taken to be $\lambda_{max} \approx 0.55\mu\text{m}$ and no response again for $\lambda > \lambda_{max}$. This appears to be a reasonable zeroth approximation to phototube response [9] leading to the requirement that about 1,000 photons have to be emitted for $\lambda_{min} < \lambda < \lambda_{max}$. We integrate eq. (5) for a fixed $\lambda_{max} = 550\mu\text{m}$ and for several values of λ_{min} , using an empirical formula [9] for the frequency dependence of the index of refraction of water. This formula is a fair approximation to the frequency dependence found in ref. [10]. The empirical formula reads:

$$\epsilon = 1.76253 - 0.0133998 (\lambda)^2 + \frac{0.00630957}{(\lambda)^2 - 0.0158806}, \quad (6)$$

the wavelength λ being measured in μm . Using (6) one has to calculate the integral:

$$F = \int_{\lambda_{min}}^{\lambda_{max}} d\lambda \frac{1}{\epsilon \lambda^2} (\epsilon - 1) \frac{d}{d\lambda} \frac{1}{\sqrt{\epsilon} \lambda} \quad (7)$$

The integration over frequencies was replaced by an integration over wavelengths, using the relation $\omega \lambda = 2\pi \epsilon^{-1/2}$. The integral F is plotted as a function of λ_{min} in Fig. 1, while keeping λ_{max} fixed. It is to be noted that in contrast to Cherenkov




Figure 1: The function F in eq. 7 plotted as a function of λ_{min} , keeping λ_{max} at $550\mu\text{m}$. Units: F is plotted in μm^{-3} and λ in μm .

radiation by a charge, the emission rate grows as ω^2 , thus magnetic Cherenkov radiation is more sensitive to detector response in the ultraviolet.

In the following table we list values of the energy, E_{1000} , necessary to generate 1,000 Cherenkov photons over a distance of 1 km, (typical of the size of future detectors) for some relevant numbers of compactified dimensions. Because of the

Table 1: E_{1000}/M_* needed for the emission of 1,000 Cherenkov photons/km

n	2	3	4	5	6	7	10	14	17	20
$^{10}\log(E_{1000}/M_*)$	10	6.88	5.33	4.41	3.8	3.37	2.62	2.14	1.99	1.84

structure of eq. (5), the energies listed are measured in units of M_* . It is to be noted that the table above can only serve as a benchmark. In any present and future detector, the minimal number of photons required depends on a geometrical factor: the fraction of the detector volume or surface covered by PMTs, depending on the specific detector type. Since the coverage is never 100%, the minimal number of photons is higher than exhibited above. However, based on the data presented, it is easily calculable for any specific detector.

So far, we have not taken into account an essential feature of the theory as described here. *In order for the usual theory of Cherenkov radiation be valid, one has to make sure that the target medium responds collectively and not as a collection of individual atoms.* One can establish a criterion for a collective response rather easily.

Assuming that the medium responds as a macroscopic body, one finds that it can take up momentum, but not energy. As a consequence, the equation of energy conservation reads:

$$p_i \approx E_i = \sqrt{p_f^2 + M_{KK}^2} + \omega. \quad (8)$$

In eq. (8), ω stands for the energy of the emitted photon, p_i, p_f for the initial and final momenta of the neutrino, respectively and M_{KK} for the mass of the KK excitation. On noting that ω is typically in the visible or in the near-UV, one finds that its contribution can be safely neglected in eq. (8). Thus, from eq. (8), one easily finds an approximate expression of the momentum transferred to the target medium, *viz*

$$q = p_i - p_f \approx \frac{M_{KK}^2}{2E_i} \quad (9)$$

The medium responds collectively if $1/q \geq d$, where d is the average intermolecular distance. In water $d \approx 10^{-8}$ cm; the corresponding energy being $1/d \approx 2$ keV. Very little is known about water or ice under extreme conditions, such as the ones prevailing at the sites of NESTOR or AMANDA. It is expected that d is smaller there. However, we use the above value to be on the safe side. In order to obtain an estimate of the neutrino energies needed, we take $N = 1$, so that $M_{KK} \approx M_c$. It follows that the initial energy of an incident neutrino has to satisfy the approximate inequality:

$$\frac{E_i}{M_c} \geq \frac{M_c d}{2} \quad (10)$$

4 Discussion

It is clear from Table 1 that low numbers of extra dimensions are disfavored from the point of view of the observability of magnetic Cherenkov radiation. In view of the stringent lower limits on M_c given by Hannestad and Raffelt [11] it is unreasonable to expect any effect even at $E \approx 10^{20}$ eV. The situation is somewhat unclear for values of n motivated by currently existing theoretical schemes, ($n = 6, 7$), since currently there are no stringent limits comparable to the ones given by ref. [11] for a low number of compactified dimensions. Future limits on n obtained from other sources will determine whether or not magnetic Cherenkov radiation is a useful tool in the search for large compactified extra dimensions.

For now, we take the limits obtained by Anchordoqui *et al.*, ref. ([12]) as a guide, as well as our recent estimate of the value of M_* [13]. Anchordoqui *et al.* give lower limits on M_* about 1 or 2 TeV, whereas, in ref. [13] we estimate $M_* \approx 80$ TeV. The requirement of coherent response, eq. 10 depends explicitly on M_c . Given symmetric toroidal compactification, as assumed, one can calculate M_c from M_* using the ADD formula, [14]:

$$M_c = 2\pi M_* \left(\frac{M_*}{M_P} \right)^{2/n}. \quad (11)$$

Using this information, we list the minimal incident neutrino energy (in the laboratory system) assuming $n = 6$ as required by a superstring for the values of M_* mentioned above.

Table 2: Values of the compactification mass and minimal value of E_i for $n = 6$, typical values of M_*

M_* in TeV	M_c in GeV	E_i in TeV
1	0.028	0.12
2	0.069	1.2
10	0.6	90
80	9.4	22,000

Clearly, very high neutrino energies are needed in order to make magnetic Cherenkov radiation emitted by neutrinos of any flavor observable. As a consequence, the coherence requirement is not a very serious issue. However, due to its unique features, such a radiation can serve as a very useful diagnostic tool in future detectors.

Acknowledgements. We thank Peter Nemethy for useful discussions on the criteria of observability of Cherenkov radiation and for supplying the empirical formula for the index of refraction. Shmuel Nussinov kindly drew our attention to the limitations of the collective description of a continuum. We thank Haim Goldberg for pointing out an error in the original version of this paper. A.E. thanks the *Ministero dell'Istruzione, dell'Università e della Ricerca* for partial support of this research under *Cofinanziamento P.R.I.N. 2001*. G.D. and S.K.D. thank

G. Pintér for first raising the possibility of using magnetic Cherenkov radiation as a tool in the search for “new physics”.

References

- [1] N. Arkani-Hamed, S. Dimopoulos, G. Dvali and J. March-Russell, Phys. Rev. **D65**:024032 (2002).
- [2] A.E. Faraggi and M. Pospelov, Phys. Lett. **B 458**, 237 (1999).
- [3] G.C. McLaughlin and J.N. Ng, Phys. Lett. **B 470**, 157 (1999).
- [4] K. Agashe and G.-H. Wu, Phys. Lett. **B 498**, 230 (2001).
- [5] R.F. Sawyer, Phys. Rev. **D 46**, 1180 (1992).
- [6] A.N. Ionissian and G.G. Raffelt, Phys. Rev. **D55**: 7038 (1997).
- [7] V.L. Ginzburg, Zh. Eksp. Theor. Fiz. **10**, 608 (1940). (This paper is in Russian; its results are quoted by various authors; however, to our knowledge, it has never been translated in its entirety.)
- [8] <http://pdg.lbl.gov>
- [9] P. Nemethy, private communication.
- [10] I. Thormahlen, J. Straub and U. Grigul, J. Phys. Chem. Ref. Data, **14**, 933 (1985).
- [11] S. Hannestad and G.G. Raffelt, Phys.Rev. Lett. **88**:071301 (2002)
- [12] L.A. Anchordoqui, J.L. Feng, H. Goldberg and A.D. Shapere, Phys. Rev. **D66**: 103002 (2002).
- [13] William S. Burgett, G. Domokos and S. Kovesi-Domokos, hep-ph/0209162. (Paper presented at ICHEP 2002, Amsterdam.)
- [14] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B429**, 263 (1998).